

2

OFFICE OF NAVAL RESEARCH

AD-A252 091



Contract N00014-90-J-1902

R&T Code 400x070yip

Technical Report No. 1

Stereoselective Oxidation of an Eleven-Membered Heterocycle

by

Paul B. Savage, John M. Desper and Samuel H. Gellman

DTIC
ELECTE
JUN 03 1992
S A D

Tetrahedron Letters, in press

Department of Chemistry
University of Wisconsin
1101 University Ave.
Madison, WI 53706

January 1992

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.


92 5 29 024

92-14253



REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a REPORT SECURITY CLASSIFICATION Unclassified			1b RESTRICTIVE MARKINGS		
2a SECURITY CLASSIFICATION AUTHORITY			3 DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release and sale; distribution unlimited.		
2b DECLASSIFICATION / DOWNGRADING SCHEDULE					
4 PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 1			5 MONITORING ORGANIZATION REPORT NUMBER(S)		
6a NAME OF PERFORMING ORGANIZATION University of Wisconsin		6b OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION Office of Naval Research	
6c. ADDRESS (City, State, and ZIP Code) Department of Chemistry, Univ. of Wisconsin 1101 University Avenue, Madison, WI 53706			7b. ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable) ONR		9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000			10 SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO 61152N	PROJECT NO 400 X	WORK UNIT ACCESSION NO 400Y070
11. TITLE (Include Security Classification) Stereoselective Oxidation of an Eleven-Membered Heterocycle					
12 PERSONAL AUTHOR(S) P.B. Savage, J.M. Desper, S.H. Gellman					
13a TYPE OF REPORT Technical		13b TIME COVERED FROM _____ TO _____		14 DATE OF REPORT (Year, Month, Day) 6 February 1992	
15 PAGE COUNT					
16 SUPPLEMENTARY NOTATION To be published in Tetrahedron Letters					
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
07	03				
19 ABSTRACT (Continue on reverse if necessary and identify by block number) <p>Oxidation of macrocyclic phosphine oxide-dithioether 1, with either mCPBA or H_2O_2/AcOH, produces only dl-2. Crystallographic and VT-^{13}C NMR data suggest that this selectivity arises from the conformational preference of 1.</p> <div style="text-align: center;">  <p>1 dl-2</p> </div>					
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21 ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a NAME OF RESPONSIBLE INDIVIDUAL Harold E. Guard			22b TELEPHONE (Include Area Code) 202/696-4311 OFFICE SYMBOL ONR Code 1113		

STERESELECTIVE OXIDATION OF AN ELEVEN-MEMBERED HETEROCYCLE

Paul B. Savage, John M. Desper and Samuel H. Gellman*

S. M. McElvain Laboratory of Organic Chemistry, Department of Chemistry,
University of Wisconsin, 1101 University Ave., Madison, WI 53706

Abstract: Oxidation of macrocyclic phosphine oxide-dithioether **1** produces only the dl form of disulfoxide **2**. This selectivity can be rationalized by comparing the crystal structures of starting material and product, and on the basis of VT NMR data, which suggest that the starting macrocycle is conformationally constrained in solution.

Molecules containing multiple sulfoxide and/or phosphine oxide groups are of interest as potential hosts for cations and for neutral guests bearing multiple hydrogen bond donors.¹ As part of our exploration of such systems, we have prepared macrocycle **1** and oxidized it to disulfoxide **2**. We report here that this oxidation is selective, providing only the dl product; neither of the meso stereoisomers was detected. Structural data for starting material and product suggest that this selectivity results from the conformational preference of eleven-membered ring of **1**.

Phosphine oxide-dithioether **1** was prepared in 74% yield by allowing the corresponding dithiol² to react with 1,4-dibromobutane and K_2CO_3 in CH_3CN at room temperature. Careful oxidation with H_2O_2 in $AcOH/CH_2Cl_2$ at room temperature or with *m*-chloroperbenzoic acid in CH_2Cl_2 at $-78^\circ C$ produced only dl-**2**. In contrast, oxidation of acyclic phosphine oxide-dithioether **3** with H_2O_2 in $AcOH/CH_2Cl_2$ produced a mixture of the four possible stereoisomeric forms of phosphine oxide-disulfoxide **4**. The dl-pair and the two meso forms of **4** were readily resolved into three peaks by normal phase (1% CH_3OH in CH_2Cl_2) and reverse phase (C_{18} ; 3:7 THF/ H_2O) HPLC. The dl-pair could be preparatively separated from the two meso isomers of **4** by flash chromatography³ on silica eluting with 3% CH_3OH in $CHCl_3$. Chromatographic analysis indicated that these stereoisomers were produced in similar quantities during the H_2O_2 oxidation of **3** (1.6:1:1 dl-pair/meso/meso).




Figure 1 shows the crystal structures of 1, dl-2 and dl-4. A comparison of the structures of 1 and dl-2 suggests a rationale for the observed oxidation selectivity. In the unsymmetrical conformation adopted by 1, each sulfur atom is oriented so that one lone pair of electrons is pointed into the center of the macrocyclic ring and thereby blocked from oxidation. For the pro-R sulfur atom of the structure shown for 1, the available lone pair is cis to the phosphine oxide oxygen atom, but for the pro-S sulfur atom, the available lone pair is trans to the phosphine oxide oxygen. Oxidant approach to the most accessible lone pair on each sulfur atom in the crystallographically observed conformation would generate the stereoisomeric form of 2 shown in Figure 1.⁴




Figure 1. (a) Ball-and-stick representations of (a) 1, (b) dl-2 and (c) dl-4 in the crystalline state; the hydrogen atoms have been omitted for clarity. Although not indicated here, the monosubstituted phenyl ring of dl-2 was disordered. This disorder could be accounted for as an approximately 1:1 mixture of two slightly different phenyl positions, only one of which is shown.

The hypothesis that oxidation selectivity results from steric constraints on oxidant approach requires that the eleven-membered ring of the starting material have well-defined conformational preferences under the reaction conditions.⁵ Variable-temperature proton-decoupled ^{13}C NMR data suggest that the mobility of this macrocycle is indeed restricted. At -50°C in CDCl_3 , the four carbons of the tetramethylene segment of **1** give rise to signals at 37.53, 34.37, 24.64 and 24.23 ppm. Coalescence among these resonances occurs as the temperature is raised: at $+50^\circ\text{C}$, only two signals are observed in this spectral region, at 36.36 and 25.18 ppm. Since the four CH_2 signals observed at low temperature are all of similar intensity, these VT-NMR data are consistent with the existence of a single unsymmetrical conformation that interconverts with its enantiomer rapidly on the NMR time scale only at the higher temperatures. Alternatively, these observations could result from the existence of two distinct symmetrical conformers that are in slow exchange and present in equivalent amounts at low temperature. This latter possibility is rendered unlikely (but not ruled out) by the behavior of the aromatic ^{13}C resonances. The four resonances arising from the carbons of the monosubstituted (exo-cyclic) phenyl ring and the lone resonance arising from the carbons para to phosphorus on the disubstituted phenyl rings remain sharp from -50° to $+50^\circ\text{C}$, while the rest of the aromatic signals undergo coalescence over this range. If there were indeed two non-enantiomeric conformations in slow exchange at -50°C , then all of the non-coalescing carbons would have to be accidentally equivalent in these two structures.

The crystal structures of **dl-2** and **dl-4** suggest that an interesting set of non-bonded interactions is important in determining the observed conformations. In both molecules, each of the sulfoxide oxygens is nearly eclipsing an aromatic ring C-C bond. In **dl-2**, the relevant OS-CC torsion angles are $6.3(6)^\circ$ and $8.0(5)^\circ$; in **dl-4** these angles are $15.5(7)^\circ$ and $20.2(7)^\circ$. One possible rationale for these preferences is steric: these torsion angles allow the methylene carbon attached to each sulfur atom to achieve maximum separation from the nearby aromatic ring (pertinent $\text{CH}_2\text{S-CC}$ torsion angles are $82.1(5)^\circ$ and $78.9(6)^\circ$ for **dl-2** and $87.8(7)^\circ$ and $86.9(7)^\circ$ for **dl-4**). It is also possible that these preferences result from an intramolecular attraction between the phosphine oxide oxygen atom and one or both of the sulfur atoms. A third possibility is a five-membered ring hydrogen bonding interaction between the sulfoxide oxygen and the nearby aromatic ring hydrogen atom (hydrogen positions not independently refined; O--H distances ca. 2.32-2.44 Å in **dl-2** and **dl-4**).

The steric rationale is not compelling because in the crystal structure of **1**, the $\text{CH}_2\text{S-CC}$ torsion angles ($42.5(2)^\circ$ and $49.0(2)^\circ$) are smaller than in **dl-2**, implying that some other factor induces the sulfoxide oxygen atoms to lie in or near the plane of the adjacent aromatic ring. The non-bonded PO--S distances in **dl-2** are 2.991(5) and 3.953(5) Å (O--S=O angles $162.8(2)^\circ$ and $175.8(2)^\circ$, respectively), and in **dl-4** the PO--S distances are 2.923(5) and 3.013(6) Å (O--S=O angles $171.2(3)^\circ$ and $173.2(3)^\circ$, respectively). The sum of the oxygen and sulfur van der Waals radii has been estimated to be 3.25 Å.⁶ Thus, there may be some non-bonded S--O attraction at



Availability Codes	
Dist	Avail. or Special
A-1	

work in dl-2 and dl-4, although much shorter non-bonded S--O contacts have been observed in other systems.^{6,7} A precedent for the five-membered ring S=O--H-C hydrogen bonding interaction proposed above may be found in structural data on co-crystals containing 1,3-bis(*m*-nitrophenyl)-urea reported by Etter and Panuto.⁸

In summary, our data suggest that the mobility of the eleven-membered ring of 1 is restricted, and that the macrocycle's conformational preference controls the stereochemistry of oxidation at the two sulfur atoms. Crystallographic data on phosphine oxide-disulfoxides dl-2 and dl-4 indicate that non-covalent attractions involving the S=O and P=O groups may exert conformation-directing effects at this oxidation level. The preparation of other macrocycles containing multiple S=O and P=O groups is underway in our laboratory, and the conformational and complexation properties of these molecules will be reported in due course.

Acknowledgments. We thank Professor H. J. Reich for helpful suggestions, and Mr. C. J. Rito for assistance with the VT-NMR measurements. This work was supported by the Office of Naval Research (Young Investigator Award to S.H.G.). P.B.S. is the recipient of a National Research Service Award (T32 GM08349) from the National Institute of General Medical Sciences. S.H.G. thanks the Searle Scholars Program and the American Cancer Society (Junior Faculty Research Award) for support.

References and Notes

1. For a discussion of polysulfoxide phase-transfer catalysts, see: Fujihara, H.; Imaoka, K.; Furukawa, N.; Oae, S. *J. Chem. Soc. Perkin Trans. I* **1986**, 465, and references therein.
2. Block, E.; Ofori-Okai, G.; Zubieta, J. *J. Am. Chem. Soc.* **1989**, *111*, 2327.
3. Still, W. C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, *43*, 2923.
4. It has been proposed that the stereochemistry of peracid epoxidations in chloroform solvents can be controlled by prior hydrogen bonding between the oxidant and a Lewis basic site on the substrate; for leading references, see: Kocovsky, P.; Stry, I. *J. Org. Chem.* **1990**, *55*, 3236. Although the oxygen atom of 1 should be a strong hydrogen bond acceptor, this directing mechanism does not seem to be a likely explanation for the selective formation of dl-2, because the selectivity is maintained when the oxidation is carried out with H₂O₂ in AcOH/CH₂Cl₂.
5. For leading references on product stereocontrol resulting from the conformational preferences of a macrocyclic starting material, see: (a) Still, W. C.; Novack, V. J. *J. Am. Chem. Soc.* **1984**, *106*, 1148; (b) Vedejs, E.; Gapinski, D. M. *J. Am. Chem. Soc.* **1983**, *105*, 5058; (c) Keller, T. H.; Weiler, L. *J. Am. Chem. Soc.* **1990**, *112*, 450.
6. Lynch, T. R.; Mellor, I. P.; Nyburg, S. C. *Acta Cryst.* **1971**, *B27*, 1948.
7. Kalman, A.; Parkanyi, L. *Acta Cryst.* **1980**, *B36*, 2372.
8. Etter, M. C.; Urbańczyk-Lipowska, Z.; Zia-Ebrahimi, M.; Panuto, T. W. *J. Am. Chem. Soc.* **1990**, *112*, 8415.